



## The Properties of the Inverse Trigonometric Functions Sine, Cosine, Tangent & Cotangent


In order for a function to have an inverse, the function must be one-to-one. In other words, it must pass the horizontal line test. Since the trigonometric functions are periodic, we must restrict the domain so that they will pass the horizontal line test. Therefore, the inverse functions will have a restricted range.




Domain  $[-1, 1]$   
Range  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$




Domain  $[-1, 1]$   
Range  $[0, \pi]$




Domain  $(-\infty, \infty)$   
Range  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$



Domain  $(-\infty, -1] \cup [1, \infty)$   
Range  $\left[-\frac{\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$



Domain  $(-\infty, -1] \cup [1, \infty)$   
Range  $\left[0, \frac{\pi}{2}\right) \cup \left(\frac{3\pi}{2}, \pi\right]$



Domain  $(-\infty, \infty)$   
Range  $(0, \pi)$

The **inverse sine function** is sometimes called the **arc sine**, and is abbreviated  $\arcsin(x)$  or  $\sin^{-1}(x)$ . Similarly, the other inverse functions are often called the **arc cosine** and **arc tangent**, abbreviated  $\arccos(x)$  or  $\cos^{-1}(x)$  and  $\arctan(x)$  or  $\tan^{-1}x$ .

$\sin^{-1}(x)$  is the angle in  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$  whose sine is  $x$ .

$\cos^{-1}(x)$  is the angle in  $[0, \pi]$  whose cosine is  $x$ .

$\tan^{-1}(x)$  is the angle in  $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$  whose tangent is  $x$ .

$\cot^{-1}(x)$  is the angle in  $(0, \pi)$  whose cotangent is  $x$ .

**Example:** Find the exact value of each expression without using a table or calculator.

a)  $\sin^{-1}\left(\frac{1}{2}\right)$

b)  $\arcsin\left(-\frac{\sqrt{2}}{2}\right)$

c)  $\tan^{-1}(1)$

d)  $\arccos\left(-\frac{1}{2}\right)$

### Identities

$$\csc^{-1}(x) = \sin^{-1}\left(\frac{1}{x}\right)$$

$$\sec^{-1}(x) = \cos^{-1}\left(\frac{1}{x}\right)$$

$$\cot^{-1}(x) = \begin{cases} \tan^{-1}(1/x) & \text{for } x > 0 \\ \tan^{-1}(1/x) + \pi & \text{for } x < 0 \\ \pi/2 & \text{for } x = 0 \end{cases}$$

$$\cot^{-1}(x) = \frac{\pi}{2} - \tan^{-1}(x)$$

**Example:** Find the exact value of each expression without using a table or calculator.

a)  $\arctan(\sqrt{3})$       b)  $\cot(\operatorname{arccot}(-5))$       c)  $\operatorname{arccot}\left(\frac{\sqrt{3}}{3}\right)$       d)  $\sin\left(\tan^{-1}\left(-\frac{3}{4}\right)\right)$

**Example:** Find the exact value of each composition.

a)  $\sin(\cot^{-1}(-1))$       b)  $\sin^{-1}\left(\sin\left(\frac{5\pi}{4}\right)\right)$       c)  $\arcsin\left(\cos\left(\frac{\pi}{6}\right)\right)$

d)  $\arctan\left(\tan\left(\frac{\pi}{3}\right)\right)$       e)  $\cos(\arctan(\sqrt{7}))$       f)  $\cot\left(\arccos\left(\frac{12}{13}\right)\right)$

**Example:** Find an equivalent algebraic expression for  $\sin(\arctan(x))$

**Example:** Find an equivalent algebraic expression for  $\cot(\arccos(2x))$